

## Normal deformed bands and possibility of TSD structures in the $N=92$ $^{166}\text{W}$ nucleus

S. Mukhopadhyay<sup>1,\*</sup>, S. K. Tandel<sup>2</sup>, W. C. Ma<sup>3</sup>, U. Garg<sup>4</sup>, Y. K. Gupta<sup>1</sup>, J. C. Marsh<sup>3</sup>, R. B. Yadav<sup>3</sup>, P. Premarashna<sup>3</sup>, J. T. Matta<sup>4</sup>, A. D. Ayangeakaa<sup>4</sup>, R. Chakrabarti<sup>5</sup>, R. V. F. Janssens<sup>6</sup>, M. P. Carpenter<sup>6</sup>, T. Lauritsen<sup>6</sup>, S. Zhu<sup>6</sup>, F. G. Kondev<sup>6</sup>, J. Chen<sup>6</sup>, D. J. Hartley<sup>7</sup>, C. J. Chiara<sup>8</sup>, and C. Petrache<sup>9</sup>

<sup>1</sup>Nuclear Physics Division, Bhabha Atomic Research Centre, Mumbai 400085, India

<sup>2</sup>UM-DAE Centre for Excellence in Basic Sciences, Mumbai 400098, India

<sup>3</sup>Department of Physics, Mississippi State University, Mississippi State, Mississippi 39762, USA

<sup>4</sup>Physics Department, University of Notre Dame, Notre Dame, Indiana 46556, USA

<sup>5</sup>Department of Physics, University of Mumbai, Vidyanagari, Mumbai 400098, India

<sup>6</sup>Physics Division, Argonne National Laboratory, Argonne, Illinois 60439, USA

<sup>7</sup>Department of Physics, USNA, Annapolis, Maryland 21402, USA

<sup>8</sup>Department of Chemistry and Biochemistry, University of Maryland, College Park, Maryland 20742, USA and

<sup>9</sup>CSNSM and CNRS/IN2P3, F-91405 Orsay, France

### Introduction

Triaxial Strongly Deformed (TSD) band structures have been observed in several Lu-Hf-Ta isotopes in the  $A\sim 165$  region. The occurrence of TSD bands in Lu ( $Z=71$ ) and Hf ( $Z=72$ ) isotopes is not consistent with predicted proton and neutron shell gaps, probably due to inadequate knowledge of energies of high- $j$  orbitals at large triaxiality. Based on calculated shell gaps, among the Lu isotopes,  $^{165}\text{Lu}_{94}$  and  $^{168}\text{Lu}_{97}$  are good candidates for TSD structures. However, the strongest TSD band in  $^{163}\text{Lu}_{92}$  is observed to be three times more strongly populated than the corresponding one in  $^{165}\text{Lu}_{94}$ , and TSD structures have not yet been observed in  $^{168}\text{Lu}_{97}$ . Further, no TSD bands have been observed in  $^{166}\text{Hf}_{94}$ , which is located at the centre of the predicted TSD island. The above observations appear to indicate that, at large triaxiality, TSD structures may be favored at  $N=92$  than at  $N=94$ .

Calculations by R. Bengtsson indicate that TSD band in  $^{166}\text{W}$ , with a positive  $\gamma$  value should become yrast at spin,  $I\sim 44\hbar$ . The aforesaid systematic calculations of TSD and

Normal Deformed (ND) structures in Hf, Yb and W isotopes further suggest that the  $N=92$  isotones in these three elements have higher excitation energies at normal deformation in the spin range of  $30\text{--}50\hbar$  [1]. So, the favorable position of TSD bands in the  $N=92$  isotones may be an outcome of the fact that the band at normal deformation goes up in energy for this neutron number.

There were two previous measurements to study high-spin states in  $^{166}\text{W}$ . In the first measurement, a  $^{28}\text{Si}$  beam was used by Gerl *et al.* [2] and four HPGe detectors were used to record data. In the second measurement by Simpson *et al.* [3], the  $p2n$  reaction channel was used, and the deexciting  $\gamma$  rays were detected using the POLYTESSA array. The positive-parity yrast band was observed up to an excitation energy of 9 MeV, and spin,  $I = 30\hbar$ , well below where possible TSD bands are expected to be yrast. Two other negative-parity rotational bands were also reported up to spin,  $I = 23\hbar$ .

### Experimental details

The experiment was performed using the Argonne Tandem Linear Accelerator System (ATLAS) facility at the Argonne National Laboratory, USA, and employed a

---

\*Electronic address: somm@barc.gov.in

285 MeV  $^{60}\text{Ni}$  beam to populate high-spin states in  $^{166}\text{W}$  with the  $^{110}\text{Pd}(^{60}\text{Ni},4n)$  reaction. Two self-supporting  $^{110}\text{Pd}$  foils of thickness  $\sim 540 \mu\text{g}/\text{cm}^2$  were stacked together and used as targets. The emitted  $\gamma$  rays were detected with the Gammasphere array which, at the time of the experiment, comprised of 82 Compton-suppressed high-purity Ge detectors. Beam wobbling was employed during the measurement, and a total of  $2.2 \times 10^9$  three- and higher-fold coincidence events were recorded for further analysis.

## Results and Discussion

The data analysis has revealed seven new high-spin rotational band structures and more than one hundred new  $\gamma$  transitions. The band built on the ground state has been extended up to an excitation energy,  $E_x = 14.4 \text{ MeV}$  and spin,  $I = 40 \hbar$ . The other two previously known negative-parity rotational bands have also been extended (Fig. 1) up to  $I = 41 \hbar$  and  $I = 38 \hbar$  (bands 1 and 2 in Ref. [3], respectively). An analysis of DCO ratios was performed to determine multiplicities of the  $\gamma$  rays and spins of excited levels were correspondingly assigned.

Cranking calculations using the Ultimate Cranker code have been performed for  $^{166}\text{W}$  and these indicate that the yrast positive-parity band built on the ground state has a prolate collective shape ( $\epsilon_2 \approx 0.17$  and  $\gamma \approx 0^\circ$ ). The first rotation alignment in this band can be attributed to the neutron  $i_{13/2}$   $AB$  crossing at 0.26 MeV (Fig. 2). The  $AB$  crossing is blocked in the negative-parity bands with 2-quasiparticle character, including an  $i_{13/2}$  neutron, therefore the  $BC$  crossing leads to the first observed alignment at 0.34 MeV. Detailed results will be presented in the symposium.

## Acknowledgments

The authors thank the ANL operations staff at ATLAS and Gammasphere. Special thanks to J. P. Greene for target preparation. This work was supported in part by the U. S. DOE grants DE-FG02-95ER40939 (MSU), DE-AC02-06CH11357 (ANL), and U. S. NSF grant PHY-1068192 (UND).




FIG. 1: Representative spectra of negative-parity rotational bands (bands 1 and 2 in upper and lower panels, respectively). The newly observed transitions are marked with an asterisk sign.




FIG. 2: Neutron quasiparticle levels in  $^{166}\text{W}$  calculated using the Ultimate Cranker code.

## References

- [1] R. Bengtsson, *private communication*.
- [2] J. Gerl *et al.*, Nucl. Phys. A **443**, 348 (1985).
- [3] J. Simpson *et al.*, J. Phys. G: Nucl. Part. Phys. **18**, 1207 (1992).