

Structural Transitions in Hot Rotating Transitional Nuclei

R. Hemalin Subala¹ and V. Selvam²

¹Department of Physics, Immanuel Arasar College of Education,
Kanyakumari – 629 195

²Department of Physics, Rani Anna Government College for Women,
Tirunelveli - 627 008, INDIA.

The study of structural transitions of nuclei at high excitation energy and large angular momentum has led us a new phase in the study of nuclear structure. The major interest in this field is the study of evolution of nuclear shape under extreme conditions of spin and temperature. In recent times, a unified framework based on the Landau theory of phase transitions has been applied to explain the universal features of the nuclear shape transitions [1]. Due to finite number of degrees of freedom, it is necessary to include thermal shape fluctuations in order to obtain good fits to experimental observables, such as GDR built on hot nuclei [2,3]. The Landau theory offers a natural framework in which these thermal fluctuations are introduced.

The quality of Landau theory applied to transitional nuclei when the free energy is expanded up to fourth power of β is not good for lower temperatures and higher spins. Hence in heavy nuclei at medium temperatures ($T \leq 1.5\text{MeV}$), it is necessary to extend the Landau free energy up to sixth order of β [4]. We have applied this extended form of Landau theory to

study the shape evolutions of hot rotating transitional nuclei, especially for the various isotopes of Neodymium. In order to obtain the constants involved in the non-rotating component of the free energy expansion, the potential energy surface obtained by the Strutinsky procedure is used. The temperature and spin dependent moment of inertia is used, which is important in transitional nuclei and the Landau constants are extracted by fitting procedure.



In the calculations performed here the spin is varied from $I = 0 \hbar$ to $60 \hbar$, temperature is varied from 0.25 to 1.5 MeV in steps of 0.25 MeV and γ is varied from -120° to -180° in steps of -10° . In order to look for near oblate and near prolate shapes, γ in steps of -2 degrees are carried out in the region -120° to -130° and -170° to -180° .

The results obtained for example in the case of ^{154}Nd for the two temperatures $T=0.25\text{ MeV}$ and 1.5 MeV are shown in figures.

It is seen from the figures that at low temperature $T=0.25\text{MeV}$ a normal shape

transition occurs from nearly prolate to triaxial as a function of spin and the deformation increases with spin. Almost the same trend is obtained with increased deformation for the temperature $T=1.5$ MeV. It is important to note that the expansion of Landau free energy in

Landau theory of shape transitions is sufficient for obtaining superdeformed configuration in transitional nuclei at normal temperatures. The sharp shape transitions are modified due to the application of thermal fluctuations.



References:

- [1] J.J. Gaardhoje, Annu. Rev. Nucl. Part. Sci. **42**, 483-536 (1992).
- [2] Y. Alhassid, J. Zingman and S. Levit, Nucl. Phys. **A469**, 205 (1987).
- [3]. Y. Alhassid and B. Bush, Nucl. Phys. **A549**, 12 (1992).
- [4] G. Shanmugam and V. Selvam Phys. Rev. **C62**, 014302 (2000).