

Band head spin assignment of Sr isotopes of superdeformed bands

Vidya Devi^{1*} and Jagjit Singh Matharu²

¹*Institute of Engineering and Technology Baddal, Ropar, India and*

²*UIET, Punjab University, Chandigarh, India*

Introduction

Superdeformation was first proposed some thirty years ago to explain the fission isomers observed in some actinide nuclei [1]. The search for superdeformed (SD) bands in the nuclear region around mass 80 was first introduced by predictions for elongated shapes with a major-to-minor axis ratio of 2:1 in nuclei with atomic number $Z=38-40$ and neutron number $N=42-45$ [2]-[3]. Superdeformed bands (SD) have been found in $^{80-83}\text{Sr}$ [4], $^{82-84}\text{Y}$ [5], $^{83,84,86}\text{Zr}$ [6], $^{85,87}\text{Nb}$ [7], ^{88}Mo [8] and $^{89,91}\text{Tc}$ [9]. These bands are observed up to a rotational frequency of about $1.3 \text{ MeV}/\hbar$, a value twice as high as the maximum frequencies encountered for the SD bands in heavy region. Therefore the study of the different properties of SD bands left a challenging task. The SD nuclei in $A=80$ mass region are theoretically described as prolate deformed rotors with a quadrupole deformation of $\beta_2 \approx 0.5$. This study is supported by the large dynamic moments of inertia θ^2 , obtained for the SD bands in the $A=80$ mass region, which have on average a value of $\approx 25\hbar^2/\text{MeV}^1$.

Spin Assignment of SDRB's in $A < 100$ Mass Region

For the deformed nuclei the collective excitation spectrum for rotational character in even-even nuclei is

$$E(I) = A [I(I + 1)] \quad (1)$$

*Electronic address: vidya.thakur@ietbaddal.edu.in

where A is inertial parameter, hence

$$E_\gamma(I) = 4A(I - I/2) \quad (2)$$

The ratio of $E_\gamma(I)$ over spin I (EGOS) is given by

$$EGOS = \frac{E_\gamma(I)}{I - I/2} = 4A. \quad (3)$$

From energy relation of VMI model we can write

$$EGOS = \frac{E_\gamma(I)}{I - 1/2} = \frac{2}{J_0} + \frac{(I^2 - I + 1)}{CJ_0^4}. \quad (4)$$

For rigid rotor, EGOS plot is a horizontal line for correct I_0 and plots are parabolic for band head spin $I_0 \pm 2$. EGOS for a nuclei following the VMI equation of energy which will be given by three parabolic curves, and it represents the presence of second order term of Bohr-Mottelson $I(I + 1)$ expansion [10].

Result and Discussion

Figure 1-2 shows the variation of EGOS against spin I . In this figure the solid circle represent the experimental values and solid line shows the calculated values of EGOS. The EGOS for $^{84}\text{Zr}(\text{SD-1})$, $^{86}\text{Zr}(\text{SD-1})$, $^{89}\text{Tc}(\text{SD-1})$ isotopes is a horizontal line for the exact I_0 and will shift to parabola for $^{80}\text{Sr}(\text{SD-1})$, $^{81}\text{Sr}(\text{SD-1})$, $^{82}\text{Sr}(\text{SD-1})$, $^{83}\text{Sr}(\text{SD-1})$, $^{83}\text{Zr}(\text{SD-2})$, $^{90}\text{Tc}(\text{SD-1})$ when $I_0 \pm 2$ is assigned to I_0 . That is, the regular structure in the one-quasi-neutron $\nu 1/2^-$ [521] negative-parity band is attributed to the decoupling effect [10].

Figure 3 shows the variation of dynamic moment of inertia with rotational frequency $\hbar\omega$ for $^{80}\text{Sr}(\text{SD-1})$, $^{81}\text{Sr}(\text{SD-1})$ nuclei. The ^{81}Sr can be represented by $\nu 5^1\pi 5^0$ configuration, the up-bend in its dynamic moment of inertia after 1.1 MeV is mainly due to the rotational




FIG. 1: EGOS versus spin to determine the band head spin for $^{80}\text{Sr}(\text{SD-1})$ nuclei.




FIG. 2: EGOS versus spin to determine the band head spin for $^{81}\text{Sr}(\text{SD-1})$ nuclei.

alignment of a pair of protons in the $h_{11/2}$ intruder orbital i.e. top of SD band in ^{81}Sr represents $\nu 5^1\pi 5^2$ configuration [10].

Acknowledgments

The author would like to thanks IET bhadral for their facility and DAE-BRNS project for the funding.

References

[1] S. M. Polikanov, et al., Zh. Eksp. Theor. Fiz. **42** (1962) 1464; Sov. Phys. JETP 15 (1962) 1016.




FIG. 3: Dynamic moment of inertia calculated using ab fitting and Harris ω^2 expansion.

[2] W. Nazarewicz, et al., Nucl. Phys. A **435** (1985) 397.
 [3] J. Dudek, W. Nazarewicz and N. Rowley, Phys. Rev. C, **35** (1987) 1489.
 [4] C. Baktash, et al., Phys. Rev. Lett. **74** (1995) 946.
 [5] P. J. Dagnall et al., Z. Phys. A **353** (1995) 251.
 [6] D. Rudolph et al., Phys. Lett. B **389** (1996) 463.
 [7] D. R. LaFosse et al., (unpublished)
 [8] T. back et al., Eu. Phys. J. A **6** (1999) 391.
 [9] B. Cederwall et al., Eur. Phys. J. A **6** (1999) 251.
 [10] A. M. Khalaf and M. D. Okasha, Progress in Physics **10** (2014) 246.