

Microscopical approach to study K-Selection Rule violation in $K=10^+$ Isomer decay of ^{182}W

C. R. Praharaj^{1,*}, S. Bhoi², Z. Naik², S. K. Ghorui³, and S. K. Patra¹

¹*Institute of Physics, Bhubaneswar - 751005, India.*

²*School of Physics, Sambalpur University, Jyoti Vihar, Burla-768019, India. and*

³*School of Physics and Astronomy, Shanghai Jiao Tong University, Shanghai 200240, China.*

Introduction

A lot more is to be explored regarding the finite but retarded transitions of K isomers to lower K bands. So, in this paper we study the decay of K isomers and the associated K selection rule [1, 2] and its violation in deformed nuclei of ^{182}W by using a microscopic model i.e. Hartree-Fock theory.

We make a consistent application of Peierls-Yoccoz procedure [3] which gives J selection rule for reduced matrix elements for electromagnetic transitions among states of various bands, but there is no K selection rule forbidding J allowed transitions between bands.

TABLE I: J to J-2 BE(2) values in $e^2\text{fm}^4$ for $K=0^+$ and $K=10^+$ band of ^{182}W .

Transitions	BE(2;J→J-2)	Transitions	BE(2;J→J-2)
$2_{gr}^+ \rightarrow 0_{gr}^+$	0.8181×10^4	$12_{iso}^+ \rightarrow 10_{iso}^+$	0.7324×10^3
$4_{gr}^+ \rightarrow 2_{gr}^+$	0.1167×10^5	$13_{iso}^+ \rightarrow 11_{iso}^+$	0.1729×10^4
$6_{gr}^+ \rightarrow 4_{gr}^+$	0.1282×10^5	$14_{iso}^+ \rightarrow 12_{iso}^+$	0.2778×10^4
$8_{gr}^+ \rightarrow 6_{gr}^+$	0.1337×10^5	$15_{iso}^+ \rightarrow 13_{iso}^+$	0.3784×10^4
$10_{gr}^+ \rightarrow 8_{gr}^+$	0.1368×10^5	$16_{iso}^+ \rightarrow 14_{iso}^+$	0.4710×10^4

TABLE II: J to J-2 BE(2) values in $e^2\text{fm}^4$ and J to J-1 BM(1) values in μ_N^2 for $K=10^+ \rightarrow K=0^+$ band of ^{182}W .

Transitions	BE(2;J→J-2)	Transitions	BM(1;J→J-1)
$10_{iso}^+ \rightarrow 8_{gr}^+$	0.4106×10^{-4}	$10_{iso}^+ \rightarrow 10_{gr}^+$	0.3×10^{-7}
$10_{iso}^+ \rightarrow 10_{gr}^+$	0.1963×10^{-3}	$11_{iso}^+ \rightarrow 10_{gr}^+$	0.3070×10^{-6}
$11_{iso}^+ \rightarrow 10_{gr}^+$	0.2353×10^{-2}	$11_{iso}^+ \rightarrow 12_{gr}^+$	0.4614×10^{-6}
$12_{iso}^+ \rightarrow 10_{gr}^+$	0.8295×10^{-2}	$12_{iso}^+ \rightarrow 12_{gr}^+$	0.7755×10^{-5}

*Electronic address: crp@iopb.res.in

Hartree-Fock theory and Angular Momentum Projection Formalism

We start with the model space which consists of $1g_{7/2}$, $2d_{5/2}$, $2d_{3/2}$, $3s_{1/2}$, $1h_{11/2}$, $1h_{9/2}$, $2f_{7/2}$, $1i_{13/2}$, and $2g_{9/2}$ with single-particle energies -6.92, -5.30, -3.58, -3.298, -4.376, 1.0, 2.0, 3.0 and 5.5 MeV respectively for protons. Also for neutrons we have $1h_{9/2}$, $2f_{7/2}$, $2f_{5/2}$, $3p_{3/2}$, $3p_{1/2}$, $1i_{13/2}$, $1i_{11/2}$, $2g_{9/2}$, and $1j_{15/2}$ with single particle energies -10.943, -11.629, -8.407, -8.739, -7.776, -9.494, -4.049, -3.485 and -0.95. The force strength of the residual interaction (surface delta) is taken as 0.15 MeV.

The microscopic model of Hartree-Fock comprises of self consistent deformed Hartree-Fock mean field obtained with a surface delta residual interaction. To get states of good J, angular momentum projection is done by the help of projection operator [4]

$$|\psi_K^J\rangle = \frac{2J+1}{8\pi^2} \int d\Omega D_{MK}^{J*}(\Omega) R(\Omega) |\phi_K\rangle \quad (1)$$

where, $R(\Omega)$ is the rotation operator and Ω represents the Euler angles α, β, γ .

The energies and electromagnetic transition operators are calculated by finding out their matrix elements which consists of integration over Euler angles. The reduced matrix elements of tensor operator T^L with multipolarity L is expressed as

$$\begin{aligned} \langle \psi_{K_1}^{J_1} || T^L || \psi_{K_2}^{J_2} \rangle &= \frac{(J_2 + 1/2)(2J_1 + 1)^{1/2}}{(N_{K_1 K_1}^{J_1} N_{K_2 K_2}^{J_2})^{1/2}} \\ &\times \sum_{\mu(\nu)} C_{\mu\nu}^{J_2 L J_1} \int_0^\beta d\beta \sin(\beta) d_{\mu K_2}^{J_2}(\beta) \\ &\times \langle \phi_{K_1} | T_\nu^L e^{-i\beta J_y} | \phi_{K_2} \rangle \quad (2) \end{aligned}$$




FIG. 1: (a) Energy spectra of $K=0^+$ and $K=10^+$ of ^{182}W (b) Plot of J overlaps for $K=0^+$ and $K=10^+$ intrinsic states.

where,

$$N_{K_1 K_2}^J = \int_0^\pi d\beta \sin(\beta) d_{K_1 K_2}^J(\beta) \times \langle \phi_{K_1} | e^{-i\beta J_y} | \phi_{K_2} \rangle \quad (3)$$

is the overlap integral. It is to be noted that the Clebsch Gordan coefficient in Eqn. 2 for the reduced matrix element has only J selection rule and there is total absence of K-selection rule for inhibition of J allowed electromagnetic transitions.

Results and Conclusion

Angular momentum projection from $K=0^+$ and $K=10^+$ intrinsic states is done and the results of the spectra is shown in Fig.1(a) along with the overlap integral (Eqn. 3) in Fig.1(b), suggesting that the wave packets are well spread. We get the energy overlap (Eqn. 2) of the $K=10^+$ states $\langle 10_{isomer}^+ | H | 10^+ gr \rangle = 0.342 \text{ eV}$ whereas the experimental energy difference between the two states is $\Delta(10_{isomer}^+ - 10^+ gr) = 0.519 \text{ MeV}$. We find that the

interaction energy is much smaller than the energy difference (by a factor of 1.5×10^6). So, we confirm that there is no k mixing. Table I shows the BE(2) values $K=0^+$ and $K=10^+$ band. Also, the BE(2) and BM(1) for $K=10^+ \rightarrow K=0^+$ are enlisted in Table II which reveal that the transitions are finite and of the order 10^{-8} of the values of transitions within a band. It is noteworthy that there is no K selection rule in our theoretical model prohibiting E2 and M1 transition but only J selection rule. Thus, we conclude that the K selection rule violating E2 and M1 transitions from the 10^+ to 0^+ band are finite but retarded which is in agreement with experiments [1, 5] and there is no K mixing in our microscopic calculation.

Acknowledgments

We thank Prof. R. K. Bhowmik, Prof. Partha Chowdhury and Prof. S. B. Khadkikar for discussions and suggestions. CRP was partly supported by SERB grant SB/S2/HEP-06/2013.

References

- [1] F. G. Kondev, G. D. Draconlis and T. Kibedi, *At. Data and Nucl. Data Tables* **103**, 50 (2015).
- [2] A. Bohr and B. R. Mottelson, "Nuclear Structure", Volume II (World Scientific 1975).
- [3] R. E. Peierls and J. Yoccoz, *Proceedings of the Physical Society* **A70**, 381 (1957).
- [4] G. Ripka, "The Hartree-Fock Theory of Deformed light Nuclei", *Advances in Nuclear physics*, Vol 1, Baranger and E. Vogt (Plenum 1966), pp. 183-259.
- [5] S. Raman, C. W. Nestor Jr., and P. Tikkanen; *Atomic Data and Nuclear Data Tables* **78**, 1 (2001).