

L^AT_EX Determination of excitation function of (n,2n) reaction on ⁸⁷Rb from threshold to 20MeV

G. Indira^{1*} and G. Anbalagan^{2†}

¹Physics Department, Government College of Engineering, Thanjavur - 613402, INDIA and

²Department of Nuclear Physics, University of Madras, Chennai-600025, INDIA

Introduction

The evaluation of cross section data for neutron induced reaction is required for several application in nuclear energy field. The nuclear model parameter plays an essential role in determining the excitation function for a particular reaction. Also, the nuclear level density parameter is effectively involved in the prediction of nuclear reaction data. In the present work the theoretical excitation function of ⁸⁷Rb(n, 2n)^{86m}Rb reaction for the incident neutron energy ranges from 10MeV to 20MeV are determined by using TALYS1.8 nuclear reaction code.

TALYS1.8 is a computer code system which is developed for the simulation of various nuclear reaction data in the field like conventional and innovative nuclear power reactors (GEN-IV), transmutation of radioactive waste, fusion reactors, accelerator applications, medical isotope production. The input parameters in TALYS are adjusted within the acceptable limit to determine the theoretical cross-section which agreed suitably with experimental cross section, and to understand the nuclear reaction theory of the particular reaction. In this work semi-microscopic optical model potential and microscopic level density model have been used to calculate the cross section for ⁸⁷Rb(n, 2n)^{86m}Rb reaction. The calculated cross sections are compared with the EXFOR experimental data and with the other evaluated data files ENDF, TENDL-2017, CENDL 3.1, BROND 3.1, ROSFOND 2010.

Nuclear model calculation

TALYS1.8 calculation with semi-microscopic optical model potential is as described in Jeukenne-Lejeune-Mahaux(JLM) optical model potential calculation. The Mod'ele Optique Microscopique module reads the radial matter densities from the nuclear structure database and performs the folding of the Nuclear Matter (NM) optical model potential described in the literature[1] with the densities to obtain a local Optical model potential. The final NM potential for a given nuclear matter density $\rho = \rho_n - \rho_p$ and asymmetry $= (\rho_n + \rho_p)/\rho$ reads

$$U_{NM}(E)_{\rho,\alpha} = \lambda_V(E) \left[V_0(\tilde{E} \pm \lambda_{V_1} \alpha V_1(\tilde{E})) \right] + i\lambda_{W_1}(E) \left[W_0(\tilde{E} \pm (E)\alpha W_1\tilde{E}) \right] \quad (1)$$

Where E is the incident neutron energy, $\tilde{E} = E - V_c$, V_c is the coulomb field, $V_0, V_1, W_0, W_1, \lambda_V, \lambda_{V_1}, \lambda_W$ and λ_{W_1} are the real isoscalar, real isovector, imaginary isoscalar and imaginary isovector, real (isoscalar+isovector), real isovector, imaginary, and imaginary isovector normalization factors respectively. The phenomenological Optical model potential for nucleon-nucleus scattering U is defined as

$$U(r, E) = -\nu_V(r, E) - iW_V(r, E) - iW_D(r, E) + \nu_{SO}(r, E).1.\sigma + iW_{SO}(r, E).1.\sigma + \nu_c(r) \quad (2)$$

where $\nu_{V,SO}, W_{V,D,SO}$ represent the real and imaginary components of the volume(V), Central (C), spin-orbit (SO) potentials. The real part of the volume potential is

$$V_V(r, E) = V_V(E)f(r, R_V, a_V) \quad (3)$$

*Electronic address: indira@gcetj.edu.in

†Electronic address: anbu24663@yahoo.co.in

In the present calculation the parameter R_V for neutron is adjusted to 1.5

Level density

There are 6 models for level density calculations that were included in TALYS1.8. The 3 phenomenological level densities which are constant temperature Fermi gas model, Back-shifted Fermi gas model, Generalised superfluid model. The 3 microscopic level density models [2] which are the Microscopic level densities from Gorielys tables, Microscopic level densities from Hilaire's combinatorial tables, Microscopic level densities from Hilaire's combinatorial tables that includes temperature-dependent Hartree-Fock-Bogolyubov calculations using the Gogny Force [3]. The microscopic level density ρ_{HfM} can be adjusted through a constant c and pairing shift δ

$$\rho(E_x, J, \Pi) = \exp\left(c, \sqrt{E_x - \delta}\right) \times \rho_{HFM}(E_x - \delta, J, \Pi) \quad (4)$$

The constant c plays a role similar to that of the level density parameter 'a' of phenomenological models. Adjusting c and δ together gives adjustment flexibility at both low and higher energies. In the present work the microscopic level density calculations using temperature-dependent Hartree-Fock-Bogolyubov calculations using the Gogny interaction are performed to produce the theoretical calculation.

Result and Discussion

The excitation function of $^{87}\text{Rb}(n, 2n)^{86m}\text{Rb}$ reaction for the incident neutron energy ranges from 10 MeV to 20 MeV are determined by using TALYS1.8 nuclear reaction codes which is illustrated in FIG.1. The present calculation is performed by invoking Jeukenne-Lejeune-Mahaux Optical model potential, microscopic level densities temperature dependent Hartree-Fock-Bogolyubov Gogny force from Hilaire's combinatorial tables. The calculated cross-section is plotted along with the available experimental data taken from EXFOR data

library and the other evaluated data files




FIG. 1: Excitation function of $^{87}\text{Rb}(n, 2n)^{86m}\text{Rb}$, along with experimental EXFOR data and other evaluated data file

ENDF, TENDL-2017, CENDL 3.1, BROND 3.1, ROSFOND 2010. The comparison of the calculated cross-section shows good agreement with EXFOR experimental data than evaluated data files.

Acknowledgments

Authors extend their gratitude to Arjan Koning, St ephane Hilaire, Stephane Goriely who provide opportunity to work with TALYS1.8

References

- [1] E. Bauge, J.P. Delaroche, M. Girod, Phys. Rev. C 63, 024607 (2001)
- [2] A.J. Koning, S. Hilaire and S. Goriely, Global and local level density models, Nucl Phys. A810,13-76 (2008).
- [3] S. Hilaire, M. Girod, S. Goriely and A.J. Koning, Temperature dependent combinatorial level densities with the D1M Gogny force, to be published (2013).