

Sensitivity on potential parameters of fusion reaction cross section around Coulomb barrier

Neha Rani¹, Pardeep Singh^{1*}, Ravinder Kumar¹, Rajiv Kumar² and Rajesh Kharab³

¹Department of Physics, Deenbandhu Chhotu Ram University of Science and Technology, Murthal 131039, Haryana, India

²Department of Physics, Govt. P.G. College for Women, Karnal, 132001, Haryana, India

³Department of Physics, Kurukshetra University, Kurukshetra, 136119, Haryana, India

*panghal005@gmail.com

Introduction

During last few decades, the fusion dynamics of weakly bound nuclei has been investigated comprehensively with the advancement in radioactive ion beam facilities [1]. The fusion reactions involving these nuclei dominantly affected by, the diffused density distribution and low breakup threshold. The key ingredient required in fusion reactions is the interaction potential and understanding of which remains one of the major subject in nuclear physics research. However, the phenomenological Woods-Saxon potential has been used frequently to interpret the experimental results obtained through fusion reactions induced by weakly and tightly bound projectiles. Nevertheless, it is not free from the ambiguities of magnitude of potential parameters which needs proper tuning.

Therefore it is worth to investigate the sensitivity of geometrical parameters like radii and diffuseness of considered potential form while analyzing the fusion reaction data. Here, in this conference contribution we focused on the analysis of ⁶He+²⁰⁹Bi fusion reaction in the vicinity of Coulomb barrier, through Kemple version of WKB approximation and Hill-Wheeler formula, with a particular emphasis to understand the sensitivity of fusion cross section on radius diffuseness, curvature as well as on the number of partial waves.

Theoretical formalism

The total interaction potential between the colliding pair may be written as,

$$V_l(r) = V_N + V_C + V_{Ce}$$

here V_N , V_C and V_{Ce} represent nuclear, Coulomb and centrifugal parts of potential respectively. The Woods-Saxon form is used to represent the nuclear part of total interaction potential and expressed as

$$V_N(r) = \frac{V_0^N}{1 + \exp\left[\frac{(r-R)}{a}\right]}$$

V_0^N , R and a gives potential depth, range and diffuseness parameters respectively. Further the depth (V_0^N) and range (R) are parameterized as

$$V_0^N = -40 \times \frac{R_P R_T}{R_P + R_T} \quad (1)$$

$$\text{with } R_{P(T)} = r_0(A_{P(T)})^{1/3} - 0.77(A_{P(T)})^{-1/3} \quad (2)$$

$$\text{and } R = r_0(A_P^{1/3} + A_T^{1/3}) \quad (3)$$

here R_P , R_T and A_P , A_T denote the size and mass numbers of projectile and target respectively. Now the fusion reaction cross section summed over partial waves can be evaluated through following expression [2]

$$\sigma_f = \frac{\pi}{k^2} \sum_{l=0}^{\infty} (2l+1) T_l^f \quad (4)$$

For projectiles with energy less than barrier height the transmission coefficient (T_l^f) for l -th partial wave is estimated through Kemple version of WKB approximation [3] which is written as

$$T_l^f = \frac{1}{1 + \exp\left[2 \operatorname{Im} \left(\int_{r_{in}}^{r_{out}} K_l(r) dr \right) \right]}$$

here r_{in} and r_{out} gives the inner and outer classical turning points of the potential $V_l(r)$ and are calculated with condition $V_l(r_{in/out}) = E$. The wave number ($K_l(r)$) is given by

$$K_l(r) = \frac{1}{\hbar} \sqrt{2\mu(E - V_l(r))}$$

While for projectile having energies above the barrier the analytical form of transmission coefficient for l -th partial wave has been expressed by Hill-Wheeler formula as [4]

$$T_l^f(E_{c.m.}) = \left(1 + \exp\left(\frac{2\pi}{\hbar\omega} (V_{Bl} - E_{c.m.}) \right) \right)^{-1}$$

here V_{Bl} represents barrier height for l -th partial wave while $\hbar\omega$ gives the corresponding curvature of the inverted parabola.

Results and Conclusions

In present contribution we have analyzed the degree of sensitivity of different geometrical components of total interaction potential on potential barrier and fusion reaction cross section for ⁶He+²⁰⁹Bi system in the

proximity of Coulomb barrier. We parameterized the values for potential depth and range parameter through the use of eqs. (1-3) and then tuned the diffuseness and radii parameter to reproduce the experimental value of barrier height and barrier position. For detail see ref [5].

Fig.1 depicts the sensitivity of pocket size on radius parameter (r_0), diffuseness parameter (a) of potential and angular momentum (l). It is clearly noticed from Fig.1 (A) that for a relatively large value of radius parameter, the fusion pocket sustains up to higher values of angular momentum (l) and partial waves up to $l = 60$ contributes. Since the inclusion of partial waves higher than $l=60$ vanishes the fusion pocket and may be neglected without much affecting the results [5]. Fig.1 (B) shows that the potential pocket is rather insensitive to the diffuseness parameter. It is clearly noticed from Fig.2 that fusion reaction cross section is enhanced due to increase in r_0 and a which may be attributed to the fact that increase in r_0 and a cause the lowering of barrier which shifts the incoming flux from entrance channel to fusion channel and leads to the fusion enhancement.




Fig. 1(Color online) (A) Variation in total potential for ${}^6\text{He}+{}^{209}\text{Bi}$ system with radius parameter 1.17 fm (solid line) and 1.5 fm (dashed line) for different l values with $a=0.5\text{fm}$. (B) Same as in (A) but with diffuseness 0.5fm (solid line) and 0.7 fm (dashed line) for different l values with r_0 fixed at 1.17fm.




Fig. 2(Color online) (A) Fusion excitation functions of ${}^6\text{He}+{}^{209}\text{Bi}$ system at around barrier energies corresponding to $r_0=1.17$ fm (lines) and 1.5 fm (symbols) for different l values with $a=0.5\text{fm}$. In B same as of (A) but with $a=0.5$ fm (lines) and 0.7 (symbols) for different l values with r_0 fixed at 1.17fm. The data point in (A) and (B) are taken from ref.[6]

Conclusively, the enhancement observed in fusion reaction cross section due to higher value of radii parameter over estimates the data up to large extent while the higher value of diffuseness also leads to enhancement in fusion reaction cross section which eventually reproduce the data especially in below barrier region. In nut shell, it is found that fusion reaction cross section is highly sensitive to the potential parameters and on l values.

Acknowledgement

The first author would like to acknowledge the financial support from TEQIP Phase-III.

References

- [1] L.F. Canto, P.R.S. Gomes, R. Donangelo, J. Lubian, M.S. Hussein Phys. Rep. 596, 1 (2015) (and refs. there in)
- [2] L.F. Canto, P.R.S. Gomes, R. Donangelo, M.S. Hussein, Phys. Rep. 424, 1 (2006).
- [3] E. C. Kemble, Phys. Rev. 48, 549 (1935).
- [4] D. L. Hill, J. A. Wheeler, Phys. Rev. 89, 1102 (1953).
- [5] Neha Rani, Pardeep Singh, Ravinder Kumar, Rajiv Kumar, Rajesh Kharab, Mod. Phys. Lett. A (Submitted)
- [6] J. J. Kolata et al., Phys. Rev. Lett. 81, 4580 (1998).