

## The $\alpha$ -decay half-life from resonance pole of S-matrix of a solvable potential

B. Sahu<sup>1,\*</sup>, Y. K. Gambhir<sup>2,3</sup>, S. Bhoi<sup>4</sup>, and M. Gupta<sup>3</sup>

<sup>1</sup>Department of Physics, College of Engineering and Technology, Bhubaneswar-751003, INDIA\*

<sup>2</sup>Department of Physics, IIT-Bombay, Powai, Mumbai-400076, INDIA

<sup>3</sup>Manipal Centre for Natural Sciences,

Manipal Academy of Higher Education,

Manipal-576104, Karnataka, INDIA and

<sup>4</sup>School of Physics, Sambalpur University, Burla-768019, INDIA

$\alpha$ -decay in nuclear physics (NP) studies remains of considerable topical interest specially in the production and study of very heavy nuclei in the actinide and superheavy regions. Recently, five new unstable isotopes in the actinide region have been reported. We present, here, the results of the calculations of half lives of these isotopes and the nuclides appearing in their  $\alpha$ -decay chains. As an illustration, we present and discuss the results of the decay  $^{217}\text{Ac}(Z=89) \rightarrow \alpha + ^{213}\text{Fr}(Z=87)$ . Microscopic calculations usually are carried out in the WKB framework which requires the Q-values and the  $\alpha$ +daughter potential. Due to high sensitivity of the calculated half lives on the Q-values used, usually the experimental Q-values are employed. Several phenomenological expressions based on the Viola-Seaborg formula are also available in the literature [1].

The  $\alpha$ +daughter potential can be calculated by folding the effective nucleon-nucleon interaction with the density distributions of  $\alpha$  and that of the daughter nucleus (double folding model or  $t\rho\rho$  approximation)[2]. Recently, calculations based on the S-matrix formulation have been reported [3]. There, the Schrödinger equation is solved with exact solvable potential which mimics the  $\alpha$ +daughter potential calculated in the double folding model. The resonance poles of S-matrix in the complex k-plane then yield the Q-values and half lives.

In the present S-matrix calculation, we use

the following exactly solvable  $\alpha$ +daughter potential:[4]

$$V_{eff}(r) = H_0 \left\{ 1 - \left[ \frac{1 - e^{-(r-R_0)/a}}{1 + ce^{-(r-R_0)/a}} \right]^2 \right\}. \quad (1)$$

Here, it has four parameters namely  $H_0$ ,  $R_0$ ,  $c$  and  $a$ . With the values of the radial position of the barrier obtained by using global formula  $R_0=R_B=r_0(A_1^{1/3} + A_2^{1/3})+2.72$  fm with  $r_0=1.0$  fm, the height of the barrier given by  $H_0=V_B=\frac{Z_1Z_2e^2}{R_B}(1 - \frac{a_g}{R_B})$  MeV with  $a_g=0.63$  fm along with the values of diffuseness  $a \approx 1.9589$  fm and constant  $c=0.6$ , the effective potential  $V_{eff}(r)$  (1) is shown in Fig. 1 as a solid curve. The figure clearly shows that this potential matches almost exactly with the Coulomb+nuclear potential ( shown in the same figure by dotted curve) of a typical  $\alpha$ +daughter nucleus system with mass number  $A_1=4$  and proton number  $Z_1=2$  of the projectile ( $\alpha$ ) and mass number  $A_2=213$  and proton number  $Z_2=87$  for the daughter nucleus  $^{213}\text{Fr}$  obtained by double folding model.

The s-wave radial Schrödinger equation with the above potential (1) is solved exactly to give the solution as

$$u(r) = A z^{ir} (1-z)^{-ir-is} F(p', m', n', \frac{1}{1-z}) + B z^{-ir} (1-z)^{ir+is} F(p, m, n, \frac{1}{1-z}),$$

where  $z = -c e^{-(r-R_0)/a}$  and  $F(p, m, n, \frac{1}{1-z})$  is the hyper-geometric function. The other terms are

$$p = \frac{1}{2} - ir - it - is, \quad m = \frac{1}{2} - ir + it - is, \\ n = 1 - 2is$$

$$p' = \frac{1}{2} + ir - it + is, \quad m' = \frac{1}{2} + ir - it + is, \\ n' = 1 + 2is,$$

\*Electronic address: bd\_sahu@yahoo.com




FIG. 1: Plot of Coulomb+nuclear potential as a function of distance for the  $\alpha+^{213}\text{Fr}$  system. The solid curve represents the results given by expression (1) with parameters stated in the text. The dotted curve stands for the double folded potential.

$r=ka$ ,  $t=[q^2(b+1)^2 - \frac{1}{4}]^{1/2}$ ,  $s=[q^2(b^2-1) + r^2]^{1/2}$ ,  $k=\sqrt{\frac{2\mu}{\hbar^2}E}$ ,  $q^2 = \frac{2\mu}{\hbar^2}H_0a^2$ ,  $b=\frac{1}{c}$ , reduced mass  $\mu=\frac{A_1A_2}{A_1+A_2}m_n$  with  $m_n$  as mass of nucleon and  $E$  stands for the incident energy.

Using the boundary condition  $u(r=0)=0$ , we get  $z_0 = -c e^{R_0/a}$  and

$$C_p = -\frac{B}{A} = \frac{z_0^{2ir}(1-z_0)^{-2ir-2is} F(p', m', n', \frac{1}{1+ce^{R_0/a}})}{F(p, m, n, \frac{1}{1+ce^{R_0/a}})}$$

As  $ce^{R_0/a} \gg 1$ ,  $1/(1+ce^{R_0/a}) \simeq 0$ ,  $F(p, m, n, 0)=1$ , we get  $C_p \simeq e^{-2\pi r} c^{-2is} e^{-2iR_0s/a}$ .

In the region  $r > R_0 = R_B$  where the potential is pure Coulombic, the Coulomb wave functions (regular  $F_0$  and irregular  $G_0$ ) for s-wave and their derivatives ( $F'_0$  and  $G'_0$ ) are considered. By requirement of continuity at  $r=R_0$ , the wave functions and their derivatives are matched at  $r=R_0$  to obtain the S-matrix denoted by  $S(k)$  as

$$S(k) = \frac{2ikF'_0 - 2iF_0f(R_0) + f(R_0)[G_0 + iF_0] - k[G'_0 + iF'_0]}{f(R_0)[G_0 + iF_0] - k[G'_0 + iF'_0]} \quad (2)$$

where  $f(R_0) = \frac{du/dr}{u} \Big|_{r=R_0}$ .

The diffuseness parameter 'a' of the potential (1) is varied around a value 2 to locate

the resonance pole  $k=k_r - ik_i$  of  $S(k)$  at the required Q-value. From this pole, we derive the decay half-life  $T_{1/2} = \frac{0.693\mu}{2\hbar k_r k_i}$ .

The effective potential (1) with parameters  $r_0 = 1.0$  fm,  $a_g = 0.63$  fm,  $c=0.6$  and  $a=1.9859$  fm having good match with the folded potential of the  $\alpha+^{213}\text{Fr}$  system is found to generate the resonance at the required energy  $Q_\alpha=9.832$  MeV giving decay half-life  $T_{1/2}^{(calc)}=6.2 \times 10^{-8}$  s. This result of calculated half-life is in good agreement with the experimental value  $T_{1/2}^{(expt)}=6.9 \times 10^{-8}$  s having 0.7 uncertainty. Similar agreement is obtained in other cases which are shown in Table 1. We conclude that the half-life calculations for  $\alpha$ -decay from the resonance poles of S-matrix of a solvable potential yield good account of the experiment.

TABLE I: Comparison between the experimental [5]  $\alpha$ -decay half-lives,  $T_{1/2}^{(expt)}$ , and calculated half-lives,  $T_{1/2}^{(calc)}$ , of different  $\alpha$ -emitters.

Parent	$E_\alpha^{(expt)}$ (MeV)	$T_{1/2}^{(expt)}$ (s)	$T_{1/2}^{(calc)}$ (s)
$^{216}\text{U}$	8.34	$3.80 \times 10^{-3}$	$4.81 \times 10^{-3}$
$^{212}\text{Th}$	7.83	$173 \times 10^{-3}$	$42 \times 10^{-3}$
$^{208}\text{Fr}$	6.61	51	21
$^{215}\text{Pa}$	8.08	$6.0 \times 10^{-3}$	$13.7 \times 10^{-3}$
$^{211}\text{Ac}$	7.481	$78 \times 10^{-3}$	$70 \times 10^{-3}$
$^{207}\text{Fr}$	6.767	1.7	5.5
$^{229}\text{Am}$	7.99	0.9	0.56
$^{225}\text{Np}$	8.63	$3.3 \times 10^{-3}$	$1.2 \times 10^{-3}$
$^{213}\text{Fr}$	6.775	16	4.1
$^{233}\text{Bk}$	7.77	21	16

## References

- [1] B Sahu, S Bhoi, *Phys. Rev. C* **93**, 044301 (2016).
- [2] Y K Gambhir, A Bhagwat, M Gupta, *Phys. Rev. C* **71**, 037301 (2005).
- [3] B Sahu, S Bhoi, *Phys. Rev. C* **96**, 044602 (2017).
- [4] T Tietz, *J. Chem. Phys.* **38**, 3036 (1963).
- [5] H M Debaraja et al., *Phys. Lett. B* **748**, 199 (2015).