

## Effect of coupling on sub-barrier fusion: The case of $^{37}\text{Cl}+^{130}\text{Te}$ system

Rudra N. Sahoo<sup>1,\*</sup>, Malika Kaushik<sup>1</sup>, Arshiya Sood<sup>1</sup>, Pawan Kumar<sup>1</sup>, Arzoo Sharma<sup>1</sup>, Swati Thakur<sup>1</sup>, Pushpendra P. Singh<sup>1,†</sup>, P. K. Raina<sup>1</sup>, Rohan Biswas<sup>2</sup>, Abhishek. Yadav<sup>2</sup>, J. Gehlot<sup>2</sup>, S. Nath<sup>2</sup>, N. Madhavan<sup>2</sup>, Md. Moin Shaikh<sup>3</sup>, Manoj K. Sharma<sup>4</sup>, B. J. Roy<sup>5</sup>, Anjali Rani<sup>6</sup>, A. Banerjee<sup>6</sup>, Unnati Gupta<sup>6</sup>, Nabendu K. Deb<sup>7</sup>, B. P. Singh<sup>8</sup>, and R. Prasad<sup>8</sup>

<sup>1</sup>Department of Physics, Indian Institute of Technology Ropar, Rupnagar-140 001, Punjab, India

<sup>2</sup>Nuclear Physics Group, Inter-University Accelerator Centre, New Delhi-110 067, India

<sup>3</sup>Variable Energy Cyclotron Centre, 1/AF, Bidhannagar, Kolkata-700064, India

<sup>4</sup>Department of Physics, S. V. College, Aligarh-202 001, U.P., India

<sup>5</sup>Nuclear Physics Division-Bhabha Atomic Research Center Mumbai-400 085, India

<sup>6</sup>Department of Physics & Astrophysics, University of Delhi-110 067, India

<sup>7</sup>Department of Physics, Gauhati University, Guwahati-781 014, Assam, India. and

<sup>8</sup>Department of Physics Aligarh Muslim University, Aligarh -202001, U.P., India

Fusion at near-to-extremely sub-barrier energies has been extensively studied in last few decades as this is useful to reveal various aspects of quantum mechanics, improve the prevailing knowledge of static and dynamic properties of nuclei, and has the probability to explore suitable condition to produce long-lived superheavy nuclei [1, 2]. In classical point of view, fusion can occur only if the system overcomes the residual barrier, formed due to equilibration of attractive nuclear potential and repulsive Coulomb potential between them. However, some experimental observations suggest that the fusion occurs also at sub-barrier energies. As per the present understanding, sub-barrier fusion occurs due to the quantum mechanical tunnelling, static and dynamic deformations, and positive Q-value neutron transfer channels (PQNT) [3, 4]. In order to understand the effect of aforesaid aspects, the fusion cross-sections of  $^{37}\text{Cl} + ^{130}\text{Te}$  system is measured from 15 % above to 10 % below the Bass barrier ( $V_B$ ).

The experiment has been carried out at Inter University Accelerator Centre (IUAC), New Delhi using the Heavy Ion Reaction Analyser (HIRA). The experimental methodology

of the present work is same with that of ref. [5]. However, a brief account of the experimental conditions is given here for ready reference. The  $^{37}\text{Cl}$  pulsed beams of bom-




FIG. 1:  $\Delta E$  - TOF spectra obtained at  $E_{c.m.}/V_B = 0.93$  and  $1.01$  where the beam-like particles are clearly separated from evaporation residues (ERs).

barding energies ( $E_{\text{Lab.}}$ ) ranging from 121 to 155 MeV were delivered from 15UD Pelletron accelerator, and bombarded on  $^{130}\text{Te}$  target, mounted inside the target chamber of HIRA and maintained  $10^{-6}$  Torr pressure. The pulsed beam was used to get a clear separation

\*Electronic address: rnsahoo@iitrpr.ac.in

†Electronic address: pps@iitrpr.ac.in

between projectile-like particles and evaporation residues (ERs), especially at below barrier energies. For  $^{37}\text{Cl} + ^{130}\text{Te}$  system, the time interval between two pulses was  $2 \mu\text{s}$ , kept slightly greater than the flight time of recoil residues ( $\approx 1.5 \mu\text{s}$  around the barrier,  $V_B \approx 134 \text{ MeV}$ ), which distinctly separate ERs and beam-like particles without overlapping the signals. The ERs are identified by making an electronic gate between TOF (subtraction of actual flight time from the time difference between two pulses) and corresponding  $\Delta E$  (energy loss by the ERs in MWPC). As a representative case, two  $\Delta E$ -TOF spectra obtained for  $^{37}\text{Cl} + ^{130}\text{Te}$  system are given in Fig.1.

TABLE I: Spectroscopic parameters of  $^{37}\text{Cl}$  and  $^{130}\text{Te}$  nuclei included in CCFULL calculations [6].

Nucleus	$E_{ex}$ (MeV)	$I^\pi$	$E(\lambda)$	$\beta$
$^{37}\text{Cl}$	1.73	vib. $(1/2)^+$	2	0.14
	3.09	vib. $(5/2)^+$	2	0.24
$^{130}\text{Te}$	0.83	rot. $2^+$	2	0.11
	2.45	rot. $4^+$	2	0.11
	1.59	vib. $2^+$	2	0.11




FIG. 2:  $\Delta E$  - TOF spectra obtained at  $E_{c.m.m.}/V_B = 0.93$  and  $1.01$  where the beam-likes are clearly separated from evaporation residues.

The fusion cross-section at different energies is estimated by using the standard expression [8]. The fusion excitation function has been obtained by plotting the fusion cross-sections

as a function of energy, and compared with the predictions of CCFULL [7] with different set of coupling as shown in Fig.2. The AW-potential parameters, *i. e.*,  $V_0 = 74.00 \text{ MeV}$ ,  $r_0 = 1.18 \text{ fm}$  and  $a_0 = 0.67 \text{ fm}$  of interacting partners are used to reproduce the measured excitation function for  $^{37}\text{Cl} + ^{130}\text{Te}$  systems. The used spectroscopic parameters in CCFULL are given in Table I.

The measured excitation function is found to be significantly enhanced relative to the one-dimensional barrier penetration model (1-DBPM). To understand this enhancement, further analysis has been performed by implementing different in-elastic excitations of interacting partners as shown in Fig.2. The coupling, *i. e.*,  $^{130}\text{Te}$  (target) : rot  $2^+$ ,  $4^+$ , vib  $2^+$  with 1 phonon, and  $^{37}\text{Cl}$  (projectile) :  $(1/2)^+$  with 2 phonon, well reproduces the observed excitation function after slightly changing the barrier by changing its radius parameter from  $r_0 = 1.18$  to  $r_0 = 1.2 \text{ fm}$ . Further data analysis is underway to understand the behaviour of fusion excitation function at sub-barrier energies. Detailed results and analysis will be presented during the Symposium.

One of the authors, R. N. S. greatly acknowledges to MHRD through IIT Ropar for financial support. Authors acknowledge the Director, IUAC for encouragement and support.

## References

- [1] A. B. Balantekin and N. Takigawa, Rev. Mod. Phys. **70**, 77 (1998).
- [2] G. Colucci *et al.*, Phys. Rev. C **97**, 044613 (2018).
- [3] A. M. Stefanini *et al.*, Phys. Rev. C **73**, 034606 (2006) and references therein.
- [4] Z. Kohley *et al.*, Phys. Rev. Lett. **107** 202701 (2011).
- [5] P. D. Shidling *et al.*, Phys. Lett. B **670**, 99 (2008).
- [6] P. Moller *et al.*, Atomic and Nuclear Data Tables **59**, 185-381(1995).
- [7] K. Hagino, N. Rowley, and T. Kruppa, Comput. Phys. Commun. **123**, 143 (1999).
- [8] Khushboo *et al.*, Phys. Rev. C **96**, 014614 (2017).