

Population of n -unbound states of ^{65}Ni via one neutron transfer reaction $^{64}\text{Ni}(^9\text{Be}, ^8\text{Be})$

Rajkumar Santra^{1,4*}, Subinit Roy¹, Haridas Pai¹, Subhendu Rajbanshi², Sajad Ali^{1,4}, Balaram Dey¹, Saikat Bhattacharjee^{1,4}, Anjali Mukherjee¹, F. S. Babra³, Md. S. R. Laskar³, Sanjoy Pal³, Rudrojyoti Palit³

¹Nuclear Physics Division, Saha Institute of Nuclear Physics, Kolkata - 700064, INDIA.

²Department of Physics, Presidency University, Kolkata-700073, INDIA.

³Tata Institute of Fundamental Research, Mumbai - 400085, INDIA.

⁴Homi Bhabha National Institute, Mumbai - 400094, INDIA.

* email: rajkumar.santra@saha.ac.in

Introduction

Transfer reaction is an indirect experimental technique to obtain the relevant quantities required to estimate the rate of astrophysical capture reaction [1]

In the present experimental investigation we explored the $1n$ transfer reaction ($^9\text{Be}, ^8\text{Be}$) on ^{64}Ni nucleus as an indirect probe for $^{64}\text{Ni}(n,\gamma)$ capture reaction. In ($^9\text{Be}, ^8\text{Be}$) reaction, the produced ^8Be quickly breaks up into two α -particles that can be detected as a clear signature of neutron transfer. The reaction $^{64}\text{Ni}(^9\text{Be}, ^8\text{Be})^{65}\text{Ni}$ has a Q -value of 4.43 MeV. Being a positive Q -value reaction, the population probability of states above the n -threshold ($S_n=6.098$ MeV) is high. We attempted the detection of γ -rays in coincidence with reaction α -particle for a high resolution determination of level energies and γ -branching factor of residual ^{65}Ni nucleus.

The branching factor will be used subsequently in $^{64}\text{Ni}(n, \gamma)$ capture reaction. The capture reaction $^{64}\text{Ni}(n, \gamma)$ has smallest Maxwellian Averaged Cross Section (MACS) among the even-even Ni-isotopes[2] and may act as a bottleneck in the formation of ^{65}Cu and other heavier nuclei in the s -process nucleosynthesis chain.

Experimental details and analysis

The experiment was performed using ^9Be (30 MeV) beam (current~5 nA) from Pelletron Linac Facility (PLF) in Mumbai. A self-supporting foil of ^{64}Ni ($\sim 500 \mu\text{g}/\text{cm}^2$) was used as the target. To detect outgoing ^8Be from $1n$ -

transfer reaction, we used CsI(Tl) detector for charged particle detection. The detector were put on both sides of the beam line covering an angular region from 22° to 67° in the reaction plane. CsI(Tl) detectors, each of size $15 \times 15 \text{ mm}^2$, were placed approximately 5cm away from the target center on each side of the beam axis. Tantalum absorbers of thickness $30 \text{ mg}/\text{cm}^2$ were used before the scintillator detectors to stop the elastically scattered particles from entering the detectors. De-exciting γ -rays of residual nuclei were detected using the γ -detector setup consisting of 14 Compton-suppressed Clover detectors placed at $40^\circ, 90^\circ, 140^\circ, 115^\circ$ and 157° with respect to the beam direction. Data were recorded in list mode in a digital data acquisition system (DDAQ) based on Pixie-16 modules of XIA-LLC, which provides both energy and timing information. The γ -ray data were sorted using Multiparameter time stamped based Coincidence Search (MARCOS) [3] program to generate one dimensional histograms, γ - γ matrix, and γ - γ - γ cube for offline analysis. RADWARE software package [4] were used for subsequent analysis.

Results and Discussions

In Fig. 1, two representative γ -spectra of residual ^{65}Ni nucleus produced in the $1n$ -transfer channel have been shown. Some of known γ -lines like 310.4 keV, 382.5 keV and 1610.4 keV are marked. Decay γ -lines from resonance states in ^{65}Ni beyond the n -threshold are shown in Fig. 2. The direct transitions to the ground state of ^{65}Ni are marked in the figure. However, the final confirmation of these states will be established by gating with the α -spectrum from CsI(Tl)

detector data. A representative 2D particle spectrum is shown in Fig. 3. Attempt will be made to identify the 2α or ${}^8\text{Be}$ band in single CsI(Tl) detector spectrum [5] to distinctly identify the γ -lines of ${}^{65}\text{Ni}$ nucleus through gating on . A detailed analysis is in progress and the results will be presented in the symposium.




Fig. 1 Gamma spectrum obtained by gating on 382.4 keV (top) and 310.4 keV (bottom) γ -rays of residual ${}^{65}\text{Ni}$ nucleus.




Fig. 2 Observed γ -rays of direct transitions from resonance states to ground state of ${}^{65}\text{Ni}$.




Fig. 3 CsI(Tl) - QDC spectrum.

Acknowledgement

The authors would like to thank the staff of Pelletron Linac Facility, Mumbai for delivering stable and uninterrupted beam during the experiment. Thanks are also due to Shri Shitalkumar V Jadhav and Dr. Balaji S Naidu for their continuous support and co-operation during the setting up of the experiment.

References

- [1] R.E. Tribble, et al., Rep. Prog. Phys. 77, 106901 (2014)
- [2] G. Cescutti, et al., arXiv:1805.01250v1 [astro-ph.SR], May 3, 2018.
- [3] R.Palit, et al., Nucl. Instrum. Methods A 680,90 (2012).
- [4] D. Radford, Nucl. Instrum. Methods A 361, 294; 306 (1995).
- [5] J.M.Almond, et al, Phys. Rev. C 86, 031307(R) (2012).