

## Mass Spectrum of Low lying Charmonium Spectrum in a Non relativistic Quark Model with Woods-Saxon Potential

A. P. Monteiro<sup>1</sup>, Praveen P D'Souza<sup>1 2,\*</sup>, Navya G R<sup>1</sup>, and K. B. Vijaya Kumar<sup>2</sup>

<sup>1</sup>*P.G Department of Physics, St Philomena College, Darbe, Puttur-574202,INDIA and*

<sup>2</sup>*Department Physics, Mangalore University, Mangalagangothri P.O., Mangalore - 574199, INDIA*

### Introduction

Charmonium system is a powerful tool for the study of forces between quarks in QCD in non perturbative regime. Studies of charmonia production can improve our understanding of heavy quark production and the formation of bound states. Our work is based on a non-relativistic potential model framework which studies the mass spectra. The potential model works well in describing the heavy quarkonia states, especially for the charmonium states below the open-charm threshold. However, above this threshold, there are still many predicted states that have not been observed yet. In recent years, charmonium physics has gained renewed strong interest from both the theoretical and the experimental side, due to the observation of charmonium-like states, such as  $Y(4260)$ ,  $Y(4360)$ , and  $Y(4660)$ [2]. A narrow resonance  $X(3872)$  was first observed in exclusive B decays  $B^\pm \rightarrow K^\pm X(3872)$ ,  $X(3872) \rightarrow \pi^+\pi^-J/\psi$  by Belle collaboration, with mass  $M = 3872.0 \pm 0.6(\text{stat}) \pm 0.5(\text{sys})$  MeV. The CDF II collaboration, D0 collaboration and BaBar collaboration.  $Y(3940)$  was discovered in exclusive B decays  $B \rightarrow KY(3940)$ ,  $Y(3940) \rightarrow \omega J/\psi$  by Belle collaboration.  $X(3940)$  was observed by Belle collaboration in  $e^+e^- \rightarrow J/\psi X(3940)$ ,  $X(3940) \rightarrow D^*\bar{D}$  with mass  $M = 3943 \pm 6 \pm 26$  MeV.  $Y(4260)$  was first observed by BaBar collaboration in initial state radiation events  $e^+e^- \rightarrow \gamma_{\text{ISR}}Y(4260)$ ,  $Y(4260) \rightarrow \pi^+\pi^-J/\psi$  with mass  $M \sim 4260$  MeV. This state was confirmed by CLEO collaboration. Other decay channels such as  $Y(4260) \rightarrow \pi^0\pi^0J/\psi$  and  $Y(4260) \rightarrow K^+K^-J/\psi$  have also been ob-

served by CLEO collaboration. Many other experimentally observed exotic states such as  $X(3915)$ ,  $Z(3930)$ , etc have lead to the deeper understanding of the charmonium physics and unravelled many mysteries[1]

### Theoretical Background

In a potential model approach the entire dynamics of quarks in a meson is governed by a Hamiltonian has kinetic energy term ( $K$ ) and a potential energy ( $V$ ), that is [3],

$$H = K + V.$$

$$K = M + \frac{p^2}{2\mu}$$

Here  $p$  is the relative momentum,  $\mu = \frac{m_Q m_{\bar{Q}}}{m_Q + m_{\bar{Q}}}$  is the reduced mass of the  $Q\bar{Q}$  system, where  $m_Q$  and  $m_{\bar{Q}}$  are the masses of the individual quark and anti quark respectively and  $M$  is the total mass of quark and antiquark [3]. In most of the cases quark-antiquark potentials ( $V(r)$ ) are reasonably considered in order to correlate results with experiment. Quark-antiquark potential may be of the form;

$$V(r) = V_{ws} + V_{CONF}(r) + V_0 \quad (1)$$

the woods saxon potential  $V_{ws}$  defined as,

$$V_{WS} = \frac{-v_0}{1 - \exp(\frac{r-R}{a})} \quad (2)$$

where  $r$  is the radial separation between the two heavy quarks and  $\alpha_s$  is the strong coupling constant and  $V_0$  is constant. For our model we have chosen the linear potential which represents the non perturbative effect of QCD that confines quarks within the color singlet system [4].

\*Electronic address: praviphysics@gmail.com

where  $a_c$  is the confinement strength and  $\lambda_i$  and  $\lambda_j$  are the generators of the color SU(3) group for the  $i^{th}$  and  $j^{th}$  quarks. It should be noted that the two body confinement potential, has symmetric and antisymmetric terms. In order to obtain the  $Q\bar{Q}$  spectrum, we have solved the Shrodinger equation with Hamiltonian equation using the variational method:

$$E(\psi) = \frac{\langle \psi | H | \psi \rangle}{\langle \psi | \psi \rangle} = \langle H \rangle$$

In our work, we have used the three-dimensional harmonic oscillator wave function which has been extensively used in atomic and nuclear physics is used as the trial wave function for obtaining the  $Q\bar{Q}$  mass spectrum.

$$\psi_{nlm}(r, \theta, \phi) = N \left(\frac{r}{b}\right)^l L_n^{l+1/2}\left(\frac{r}{b}\right) \exp\left(-\frac{r^2}{2b^2}\right) Y_{lm}(\theta, \phi)$$

where  $|N|$  is the normalizing constant given by

$$|N|^2 = \frac{2\alpha^3 n!}{\sqrt{\pi}} \frac{2^{(2(n+l)+1)}}{(2n+2l+1)!} (n+l)! \quad (4)$$

and  $L_n^{l+1/2}(x)$  are the associated Laguerre polynomials,

### Results and Discussions

There are five parameters in our model. These are the the mass of the bottom quark  $M_b$ , the confinement strength  $a_c$ , the quark-gluon coupling constant  $\alpha_s$ , potential depth  $V_0$ ,  $R$  and  $a$  [7]. These parameters are adjusted until the energy of the trial wavefunction is minimized. The resulting trial wavefunction and its corresponding energy are variational method approximations to the exact wavefunction and energy. Within the framework of the non-relativistic potential model, the trial wave function of the heavy quarkonium satisfies the Schrodinger equation. The variational method has extensively been used and seems to be successful in many aspects-

$$V_0 \approx 50 \text{ MeV}; a \approx 0.6 \text{ fm}; R = 8 \text{ fm}; M_b = 4.645 \text{ GeV};$$

$$a_c = 135.0 \text{ MeV fm}^{-1}$$

Table 1 The Mass Spectrum of Charmonim (MeV)

$n^{2S+1}L_J$	The Mass	$M_{exp}$ MeV	[2] MeV
$1^1S_0$	2982.53	$2983.60 \pm 0.7$	2990.40
$2^1S_0$	3635.37	$3639.2 \pm 0.11$	3646.50
$3^1S_0$	3963.60	....	4071.90
$4^1S_0$	4353.60	....	....
$5^1S_0$	4586.42	....	....
$1^3S_1$	3094.54	$3096.916 \pm 0.011$	3085.10
$2^3S_1$	3684.38	$3686.108 \pm 0.018$	3682.10
$3^3S_1$	4040.62	$4039 \pm 1$	4100.20
$4^3S_1$	4453.00	....	....
$5^3S_1$	4473.29	....	....

even though various forms of  $q\bar{q}$  potentials have been predicted, potential model approach is still a powerful tool for investigating the properties of quarkonia owing to its great predictive power and mathematical simplicity. Also, the potential model provides a framework to understand the structure of the spin dependent interaction in QCD, the form of confinement, Coupled channel effects decay properties etc.

### References

- [1] C. P. et al., Chinese Physics C 40 (2016) 100001
- [2] Jielei Zhang and Rumin Wang. Advances in High Energy Physics 2018 (5),
- [3] Bhaghyesh and K. B. Vijaya Kumar, International Journal of Modern Physics A 27 (2012) 1250127.
- [4] Antony Prakash Monteiro, Manjunath Bhat and K. B. Vijaya Kuma, International Journal of Modern Physics A **32**, 04,(2017)
- [5] Antony Prakash Monteiro, Manjunath Bhat and K. B. Vijaya Kumar. Physics Review D, 054016 **95** (2017)
- [6] Jun-Zhang Wang, Zhi-Feng Sun, Xiang Liu, Takayuki Matsuki arXiv.org :arXiv:1802.04938
- [7] Proceedings of the DAE-BRNS Symp. on Nucl. Phys. 60 (2015)