

## Intermittency analysis of generated charged particles in central Pb-Pb collisions using EPOS3 model

Salman Khurshid Malik\*  
 Department of Physics,  
 University of Jammu, Jammu-180006, INDIA

### Introduction

One of the primary goals in investigating the collisions of heavy nuclei at relativistic energies is to understand the nature of phase-transition from hadronic matter to quark-gluon plasma and vice versa. The collision experiments give us an opportunity to test the predictions of QCD and also to comprehend other processes involved. A system in the state of QGP is expected to undergo quark-hadron phase transition with the expanding and cooling. Although, a number of signals have been proposed, but still a little is known.

The multiplicity fluctuations of hadrons produced in heavy-ion collisions can indicate a phase transition of QGP and hence, study of fluctuations can be used as a signal. The reason is that the fluctuations are substantial near the critical point for statistical systems. A lot of effort has been made in studying large-density fluctuations in hadronic and heavy-ion collisions, which might give us an insight into the phase-transition.

One of the most suitable methods to identify the existence of non-statistical fluctuations is the Normalized Factorial Moments (NFMs) [1]. This method has a scaling nature with diminishing phase-space domains, known as Intermittency, which is mainly an arrangement with the short-range correlations among the particles produced in the heavy ion collisions at high energies. Intermittency may be due to the self-similarity in the random cascading of elementary processes or it may be because of the self-similarity in nuclear collisions by the second-order phase transition

from QGP to the hadronic state.

In the present work, intermittency in two dimensional phase-space of kinematical variable pseudorapidity ( $\eta$ ) and azimuthal angle ( $\phi$ ) of the events generated using EPOS3 [2] model for Pb-Pb collisions at  $\sqrt{s_{NN}}=2.76$  TeV will be presented.

### Methodology

The normalised factorial moments,  $F_q$  of order  $q$ , in its vertical form are defined as [1]:

$$F_q(M) = \left\langle \frac{1}{M} \sum_{m=1}^M \frac{f_q^{ev}(M)}{\langle N \rangle^q} \right\rangle \quad (1)$$

$\langle \dots \rangle$  is for averaging over all events ( $N_{events}$ ) with,

$$f_q^{ev}(M) = n_m(n_m - 1) \dots (n_m - q + 1) \quad (2)$$

where  $n_m$  is the number of charged particles in the  $m^{th}$  bin ( $m=1,2,3,\dots,M$ ) and  $\langle N \rangle$  is the mean multiplicity of hadrons in the pseudorapidity interval  $\Delta\eta (= \eta_{max} - \eta_{min})$ , which in turn is divided into  $M$  bins of equal width,  $\delta\eta = \frac{\Delta\eta}{M}$ .

It has been proved [1] that in case of statistical fluctuations, for a smooth pseudorapidity distribution,  $\langle F_q \rangle$  is primarily independent of the bin width  $\delta\eta$  in the limit  $\delta\eta \rightarrow 0$ . In the limit of small bin size and when the fluctuations are dynamical in nature, the following proportionality holds:

$$\langle F_q \rangle \propto (M)^{\varphi_q}; \delta\eta \rightarrow 0 \quad (3)$$

This power law behaviour is known as intermittency and  $\varphi_q$ , intermittency exponent measures the strength of fluctuations.

It is observed [5] that for the second-order phase transition, the power law behaviour is:

$$F_q \propto F_2^{\beta_q} \quad (4)$$

---

\*Electronic address: maroozmalik@gmail.com

where

$$\beta_q = (q-1)^\nu, \nu = 1.304 \quad (5)$$

$\nu$  (the scaling exponent) is significant in a way that it can be used to characterize scaling properties of multiparticle production. It is also independent of the details of the system, i.e., it is universal.

After observing the power law and self-similarity, the analysis in terms of fractal parameters [3] is crucial. The strength of intermittency effect is related to generalized dimensions by :

$$D_q = 1 - \frac{\varphi_q}{(q-1)} \quad (6)$$

$$\text{or } D_q = 1 - d_q, d_q = \frac{\varphi_q}{q-1} \quad (7)$$

where  $d_q$  are the anomalous fractal dimensions of fractals and multifractals in terms of intermittency exponents,  $\varphi_q$

Although, data has been interpreted in terms of anomalous dimensions, the relationship between intermittency and fractality or multifractality is still vague. *Bialas* and *Hwa* [4] suggested that if a second-order phase transition takes place from QGP to a hadron phase in the thermodynamic equilibrium, then the produced particles will show intermittency with  $d_q$  independent of  $q$ .

## Observations

EPOS3 events are studied for the intermittency and the dependence of power-law slopes will be investigated. Variation of the Anomalous fractal dimensions ( $d_q$ ), Generalised fractal dimensions ( $D_q$ ) and their variations with  $q$  will also be presented.

## Summary

Charged particle multiplicity distributions for Pb-Pb collisions at  $\sqrt{s_{NN}} = 2.76$  TeV using EPOS3 model are studied for their local

multiplicity fluctuations.  $\nu$  the scaling index value obtained for the model is the characteristic of the generator.




FIG. 1: Pseudorapidity distribution of the EPOS3 event sample compared with ATLAS data for Pb-Pb collisions at  $\sqrt{s_{NN}} = 2.76$  TeV.

## Acknowledgments

The heartfelt thanks are due to *Dr. R.C. Hwa* for his valuable discussions. I also acknowledge the timely and quick help provided by *Tanguy Pierog* and *Klaus Werner* for the installation and setting up of EPOS3 model.

## References

- [1] A. Bialas and R. Peschanski, Nucl. Phys. B **273**, 703 (1986); B **308**, 867 (1988).
- [2] T. Pierog, I. Karpenko, J.M. Katzy, E. Yatsenko, and K. Werner, Phys. Rev. C **92** (2015) 034906
- [3] P Lipa and Buschbeck, Phys. Lett. B **223**,465 (1989);
- [4] A. Bialas and R.C. Hwa, Phys. Lett. B **253**,436 (1991);
- [5] R.C. Hwa and M.T. Nazirov, Phys. Rev. Lett. **69**,741 (1992);