

Measurements of neutron capture reaction relevant to r-process via Coulomb breakup of ^{32}Mg RIB

W. Sengupta,^{1,2} U. Datta,^{1,2,*} A. Rahaman,^{1,3} T. Aumann,^{4,5} S. Beceiro-Novo,⁶ K. Boretzky,⁴ C. Caesar,⁴ B. V. Carlson,⁷ W. N. Catford,⁸ S. Chakraborty,^{1,9} M. Chartier,¹⁰ D. Cortina-Gil,⁶ G. De. Angelis,¹¹ P. Diaz Fernandez,⁶ H. Emling,⁴ H. Geissel,^{4,12} D. Gonzalez-Diaz,⁴ M. Heine,^{4,13} H. Johansson,¹⁴ B. Jonson,¹⁴ N. Kalantar-Nayestanaki,¹⁵ T. Kröll,^{5,16} R. Krücken,¹⁶ J. Kurcewicz,⁴ C. Langer,⁴ T. Le Bleis,¹⁶ Y. Leifels,⁴ J. Marganiec,¹⁷ G. Münzenberg,⁴ T. Nilsson,¹⁴ C. Nociforo,⁴ V. Panin,⁴ R. Plag,⁴ R. Reifarth,⁴ M. V. Ricciardi,⁴ C. Rigollet,¹⁵ D. Rossi,^{4,16} S. Santra,^{2,18} C. Scheidenberger,^{4,12} H. Scheit,⁵ H. Simon,⁴ Y. Togano,^{17,19} S. Typel,^{4,5} Y. Utsuno,²⁰ A. Wagner,²¹ F. Wamers,⁴ H. Weick,⁴ and J. S. Winfield⁴

¹Saha Institute of Nuclear Physics, Kolkata, India

²Homi Bhabha National Institute, Mumbai, India

³Jalpaiguri Government Engineering College, Jalpaiguri, India

⁴GSI Helmholtzzentrum für Schwerionenforschung GmbH, Darmstadt, Germany

⁵Technische Universität Darmstadt, Darmstadt, Germany

⁶Universidade de Santiago de Compostela, Santiago de Compostela, Spain

⁷Instituto Tecnológico de Aeronautica, Sao Jose dos Campos, Brazil

⁸University of Surrey, Guildford, United Kingdom

⁹University of Engineering and Management, Kolkata, India

¹⁰University of Liverpool, Liverpool L69 7ZE, United Kingdom

¹¹INFN, Legnaro, Italy

¹²Physikalisches Institut, Giessen, Germany

¹³IPHC and CNRS, Université de Strasbourg, Strasbourg, France

¹⁴Fundamental Fysik, Chalmers Tekniska Högskola, Goteborg, Sweden

¹⁵Nuclear Energy Group, ESRIG, University of Groningen, Groningen, The Netherlands

¹⁶Physik Department E12, Technische Universität München, Garching, Germany

¹⁷EMMI, GSI Helmholtzzentrum für Schwerionenforschung GmbH, Darmstadt, Germany

¹⁸Nuclear Physics Division, Bhabha Atomic Research Centre, Mumbai 400085, India

¹⁹Department of Physics, Rikkyo University, Tokyo, Japan

²⁰Advanced Science Research Center, Japan Atomic Energy Agency, Tokai, Ibaraki, Japan

²¹Helmholtz-Zentrum Dresden-Rossendorf, Dresden, Germany

Introduction

The Rapid Neutron Capture or *r*-process nucleosynthesis, responsible for synthesis of $\sim 50\%$ of the heavy nuclei in the universe, is known to occur in Core-collapse Supernovae (CCSNe) and Neutron Star Mergers (NSM). There are many reaction network codes, which are used to theoretically model this *r*-process.

According to the ν -driven wind model, (n, γ) capture rate of light neutron-rich nuclei might play a role in determining the *r*-process abundance of elements [1]. But experimental measurements related to such reactions are extremely difficult owing to instability of the reactants involved. Hence, *r*-process simulations usually takes into account theoretical reaction rates from *Hauser-Feshbach* (HF) statistical model calculations, e.g. T. Rauscher calculations [2]. In the present work, we have performed experimental measurements of $^{31}\text{Mg}(n, \gamma)^{32}\text{Mg}$ capture cross-section and

*Electronic address: ushasi.dattapramanik@saha.ac.in




FIG. 1: Raw distribution of Excitation Energy, E^* of ^{32}Mg on ^{208}Pb target from Eq 1.

obtained the reaction rate through the indirect method of *Coulomb breakup* (theory discussed in [3–5]) and compared with HF calculations.

Experiment

The neutron dissociation reaction, $^{32}\text{Mg} \rightarrow ^{31}\text{Mg} + n$ is induced via Coulomb breakup of ^{32}Mg RIB off a ^{208}Pb target at $\sim 400A$ MeV. The experiment was performed at GSI, Darmstadt, Germany using the FRS-LAND setup (**S306:GSI**). The same reaction was further performed with ^{12}C as target and also in absence of any target, for background estimation. Details of experiment can be found in [6].

Data analysis

The **differential cross-section**, $d\sigma/dE^*$ of $^{31}\text{Mg} + n$ break-up channel is measured via invariant mass analysis as:

$$E^* = \sqrt{m_f^2 + m_n^2 + 2m_fm_n\gamma_f\gamma_n \times \frac{1}{(1 - \beta_f\beta_n\cos\theta)}} - M \quad (1)$$

Where, f and n refers to the fragment (^{31}Mg) and the dissociated neutron, respectively, while M is mass of the projectile, ^{32}Mg . The reaction on Pb target comprises contributions from both Coulomb (electromagnetic) and nuclear interactions. In order to eliminate said nuclear component, the data with ^{12}C as target is used and a pure Coulomb, $d\sigma/dE^*$ is obtained. Further correcting the same for instrumental *efficiency* and *acceptance*, we find the **photo-dissociation cross-**




FIG. 2: Comparison of experimental and theoretical astrophysical reaction rates.

section, $\sigma_{\gamma,n}$. The inverse **capture cross-section** $\sigma_{n,\gamma}$ is then calculated via principle of *detailed balance*, followed by the astrophysical **reaction rate**, $N_A\langle\sigma_{n,\gamma}v\rangle$. Detailed calculation scheme had been discussed in [4, 5]. The experimental reaction rate is compared with the HF calculations of TALYS code and T. Rauscher code, widely used in r-process simulations. At temperatures relevant to *strong r-process* i.e. in NSMs (~ 6 GK), the two models lie very close to our results, with a relative difference of $\sim 20\text{-}30\%$. But, the difference with theory becomes much larger in lower temperature regimes of *weak r-process* occurring in CCSNe, a trend which is also observed in [7]. Further details of the analysis and the results will be discussed in the conference.

References

- [1] M. Terasawa et al. A.P.J. **562**, 470 (2001).
- [2] T. Rauscher et al. A.D.N.D.T. **75.1**, 1 (2000).
- [3] Bertulani and Baur Phys. Rep. **163**, 299 (1988).
- [4] U. Datta, et al. Phys. Lett. B **551**, 63 (2003) & Prog. Part. Nucl. Phys. **59**, 183 (2007).
- [5] W. Sengupta et al. DAE Symp. on Nucl. Phys. **67**, C48 (2023).
- [6] A. Rahaman et al. J. Phys. G: Nucl. Part. Phys. **44**, 045101 (2017)
- [7] M. Heine et al. Phys. Rev. C **95**, 014613 (2017)