

# Hadron-hadron scattering by energy-dependent and independent interactions- a supersymmetric approach

M. Majumder<sup>1\*</sup> and U. Laha<sup>1</sup>.

<sup>1</sup>Department of Physics; National Institute of Technology, Jamshedpur, India, 831014

\*Electronic address: madhuramajumder2015@gmail.com

## Introduction:

Utilizing the theoretical underpinnings, such as Schrödinger's equation, Supersymmetric Quantum Mechanics (SQM), and the Phase Function Method (PFM) both charged and uncharged systems are investigated, with a focus on a diverse array of local and nonlocal potentials, including the intricate landscape of screened Coulomb potentials. The conversion of non-local nuclear interactions into simplified local forms addresses challenges inherent in solving the Schrödinger's equation for complex potentials.

The impact of short-range interactions between nuclei, primarily due to multi-pion exchange processes and nucleon recoil necessitates consideration of non-local nuclear interactions, represented by two variables 'r' and 'r''. Mathematical manipulation enables the conversion of multivariable non-local potentials to simplified local forms. While exact solutions exist for specific potentials, such as Coulomb, Square well, etc., others like Hulthén and Yamaguchi are solvable only for the S ( $l=0$ ) state. Various mathematical approaches, including supersymmetry factorization and Phase function methods, are employed to address this limitation, focusing on charged and uncharged systems. The  $\alpha$ - $\alpha$  system is extensively studied with different potentials to explain scattering phase shifts. The Hulthén potential is identified as an effective model for deuteron and  $\alpha$ -nucleon interactions.

The relationship between regular and irregular solutions of non-local and local potentials has been explored, emphasizing the energy dependence of the Yamaguchi potential and the need to assess if an energy-dependent potential out-performs an energy-independent one.

The Jost Function plays a central role in developing a relativistic S-matrix theory, crucial for understanding particle interactions during

scattering processes. the Jost function is instrumental in analyzing bound and resonant states, as well as low-energy scattering data. Its holomorphic nature in the upper complex k-plane ( $\text{Im } K > 0$ ) is key, as the zeros of the Jost function in this plane are directly linked to bound state energies. The On-Shell Jost Function is easily calculated from its integral representation, correlating phase values with scattering phase shifts with a  $\pi$  phase difference. For accurate modeling of few-nucleon systems (e.g.,  $\alpha$ -nucleon, p-p, n-<sup>3</sup>He), a customized centrifugal barrier term is incorporated, aligning with nuclear interaction requirements. Literature has explored diverse approaches to approximate higher partial waves.

The supersymmetry transformation operators ( $T_1, T_2, T_3, T_4$ ), integral to particle physics are exploited to create phase-equivalent potentials. The SUSY operators transition between fermionic and bosonic states, unveiling super partners in quantum field theory. They modify the initial Hamiltonian uniquely:  $T_1$  eliminates the ground state,  $T_2$  adds an extra bound state, and  $T_3$  and  $T_4$  alter the Jost Function and potential singularity. Applying these operators to a Manning-Rosen potential yields phase-equivalent potentials, and comparisons with experimental data validate the approach.

We make use of deformed Hulthén potential, delving into the derivation of regular and Jost solutions. The incorporation of energy-dependent correction terms enhances the accuracy of predictions, particularly for deuteron and triton nuclei, contributing to a nuanced understanding of nuclear forces.

## Results and Discussions:

Figures 1 to 4 depicts scattering phase shifts and cross sections data of some of the system studied




Fig.1:  $^3D_1$  energy dependent and independent phase shifts as a function of  $E_{Lab}$  with ref. [1, 2]. In Fig.1 we see phase shifts due to energy dependent potential out performs energy independent one.




Fig. 2:  $\alpha - \alpha$  scattering phase shifts for S and D waves with standard data.

The  $\alpha - \alpha$  phase shifts for S and D waves have better match with standard data for Coulomb-Nuclear-SUSY interactions, as shown in Fig. 2. Fig. 3 shows that differential scattering cross section, for  $\alpha$ -p system match well with Brockman *et al.* data. Phase-shifts generated from the phase equivalent potentials, using four SUSY transformations match well with experimental data.




Fig. 3: Differential scattering cross section for  $\alpha$ -p system at two different lab energies with standard data [3].




Fig4: Phase shifts for  $1/2(+)$ ;  $1/2(-)$  and  $5/2(+)$   $\alpha$ -n states for Phase equivalent potentials with reference data [4].

#### References:

1. R. A. Arndt, L. D. Roper, R. A. Bryan, R. B. Clark, B. J. Verwest and P. Signal, Phys Rev D **28**, 97 (1983).
2. F. Gross and A. Stadler, Phys Rev C **78**, 014005 (2008).
3. K. W. Brockman Phys. Rev. **108**, 1000 (1957).
4. G. R. Satchler, L. W. Owen, A. J. Elwin, G. L. Morgan and R. L. Walter, Nucl. Phys. A **112**, 1 (1968).